

Modulation of endoplasmic reticulum calcium pump expression during lung cancer cell differentiation

Atousa Arbabian^{1,2}, Jean-Philippe Brouland³, Ágota Apáti^{4,5}, Katalin Pászty⁶, Luca Hegedűs⁴, Ágnes Enyedi^{4,5}, Christine Chomienne^{1,2} and Béla Papp^{1,2,7,8}

1 Institut National de la Santé et de la Recherche Médicale, UMR-S 940, Paris, France

2 Institut Universitaire d'Hématologie, Université Paris Diderot, PRES Sorbonne Paris-Cité, France

3 Service d'Anatomie et Cytologie Pathologiques, Hôpital Lariboisière, Paris, France

4 Institute of Molecular Pharmacology, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Budapest, Hungary

5 Department of Molecular Cell Biology, National Blood Service, Budapest, Hungary

6 Molecular Biophysics Research Group, Hungarian Academy of Sciences and Semmelweis University, Budapest, Hungary

7 Institut National de la Santé et de la Recherche Médicale, UMR U978, Bobigny, France

8 Université Paris-13, PRES Sorbonne Paris-Cité, Bobigny, France

Keywords

calcium; differentiation; endoplasmic reticulum; lung cancer; SERCA

Correspondence

B. Papp, UMR U978 Inserm, Université Paris-13, UFR SMBH, 74 rue Marcel Cachin, 93017 Bobigny, France
Fax: +33 1 48 38 73 69
Tel: +33 6860 88309
E-mail: belapapp2@yahoo.fr

(Received 4 September 2012, revised 23 October 2012, accepted 7 November 2012)

doi:10.1111/febs.12064

Cellular calcium signaling plays important roles in several signal transduction pathways that control proliferation, differentiation and apoptosis. In epithelial cells calcium signaling is initiated mainly by calcium release from endoplasmic-reticulum-associated intracellular calcium pools. Because calcium is accumulated in the endoplasmic reticulum by sarco/endoplasmic reticulum calcium ATPases (SERCA), these enzymes play a critical role in the control of calcium-dependent cell activation, growth and survival. We investigated the modulation of SERCA expression and function in human lung adenocarcinoma cells. In addition to the ubiquitous SERCA2 enzyme, the SERCA3 isoform was also expressed at variable levels. SERCA3 expression was selectively enhanced during cell differentiation in lung cancer cells, and marked SERCA3 expression was found in fully differentiated normal bronchial epithelium. As studied by using a recombinant fluorescent calcium probe, induction of the expression of SERCA3, a lower calcium affinity pump, was associated with decreased intracellular calcium storage, whereas the amplitude of capacitative calcium influx remained unchanged. Our observations indicate that the calcium homeostasis of the endoplasmic reticulum in lung adenocarcinoma cells presents a functional defect due to decreased SERCA3 expression that is corrected during pharmacologically induced differentiation. The data presented in this work show, for the first time, that endoplasmic reticulum calcium storage is anomalous in lung cancer cells, and suggest that SERCA3 may serve as a useful new phenotypic marker for the study of lung epithelial differentiation.

Introduction

Non-small-cell lung cancer is currently the most frequent fatal malignancy worldwide with more than a million deaths occurring annually [1,2]. Although much is known about risk factors such as tobacco

smoking, or regarding molecular and genetic events that occur during lung tumorigenesis, the overall survival rate of lung cancer could not be significantly improved during the last 30 years [2]. This indicates

Abbreviations

CAG promoter, cytomegalovirus early enhancer/chicken β actin promoter; ER, endoplasmic reticulum; GFP, green fluorescent protein; SB, Sleeping Beauty; SERCA, sarco/endoplasmic reticulum calcium ATPase.

that currently available knowledge of the molecular biological behavior of lung carcinoma cells is insufficient for the development of efficient therapy. In particular, knowledge about calcium signaling in lung cancer cells is scarce, although calcium signaling is an essential component of several regulatory networks involved in the control of cell growth, differentiation and apoptosis.

Calcium-dependent cell activation is initiated by the release of calcium ions stored in the endoplasmic reticulum (ER) into the cytosol through inositol-1,4,5-*tris*phosphate receptor calcium channels [3]. In addition, calcium in the ER lumen is required for the function of various ER located chaperones such as calreticulin or calnexin involved in the conformational maturation and post-translational modification of newly synthesized proteins that transit through the organelle [4–6]. Abnormal ER calcium homeostasis can lead to cellular stress responses, growth arrest and apoptosis [4,7–9].

Sarco/endoplasmic reticulum calcium ATPase (SERCA) enzymes are essential for the dynamic equilibrium of ER calcium homeostasis. Located in the ER membrane, these enzymes sequester calcium ions from the cytosol into the ER lumen by ATP-dependent active ion transport, and thus generate a steep calcium concentration gradient between the cytosol (approximately 50–100 nM) and the ER lumen (approximately 0.1 mM) [10]. SERCA proteins are encoded by three genes (ATP2A1, 2 and 3) that can give rise to several protein isoforms by alternative splicing [11–15]. SERCA expression is tissue dependent and developmentally regulated. SERCA1 is expressed in skeletal muscle, and the SERCA2a isoform is mainly expressed in cardiomyocytes. Whereas SERCA2b is a ubiquitously expressed isoform thought to be involved in ‘housekeeping’ functions, in several cell types such as cells of hematopoietic origin, gastric and colonic epithelial cells, β -cells of islets of Langerhans or vascular endothelium, SERCA2b has been shown to be co-expressed with SERCA3 [15–26]. Because SERCA2b and SERCA3 enzymes display distinct calcium affinities [27–32], their co-expression is involved in the fine-tuning of the parameters of ER calcium sequestration according to the specific calcium storage and release requirements of the ER of a given cell type.

By controlling intra-ER calcium levels, SERCA enzymes are involved in the control of cell growth [33–36]. Small molecular SERCA inhibitors are tumor promoters [37,38], and SERCA mutations have been identified in cancer [39–43]. In this context we have previously reported that the expression of SERCA enzymes is anomalous in breast, colon as well as gastric carcinoma due to the loss of expression of

SERCA3 [18,44,45]. It has also been shown that SERCA3 expression is induced during histone-deacetylase inhibitor-induced differentiation of gastric and colon carcinoma cells [18], as well as following the inhibition of the APC/ β -catenin/TCF4 axis [44], and that normal colonic and gastric epithelium expresses SERCA3 abundantly [44]. In addition, the pharmacological modulation of SERCA activity has also been shown to enhance cell differentiation in colon carcinoma [44] and acute promyelocytic leukemia [46] cells. Taken together, these observations indicate that the functional maturation of ER calcium homeostasis that occurs during normal cell differentiation is blocked due to deficient SERCA3 expression in malignant cells, and that this phenomenon is involved in the maintenance of the neoplastic phenotype [16,47].

In order to establish whether defects of ER calcium homeostasis can be found in pulmonary neoplasia, in this work we investigated SERCA expression in normal lung *in situ* and in lung adenocarcinoma cell lines undergoing pharmacologically induced cell differentiation *in vitro*, and investigated functional calcium storage capacity by confocal microscope calcium fluorimetry using a green fluorescent protein (GFP) based calcium probe in differentiated and control cells. Our results indicate that SERCA3 is abundantly expressed in normal fully differentiated bronchial epithelium, whereas expression is highly heterogeneous in carcinoma tissue. We also show that SERCA3 expression is induced during the pharmacologically induced differentiation of lung adenocarcinoma cell lines, and that differentiation induction leads to decreased ER calcium storage.

These observations show that, similarly to other types of neoplasia, lung cancer cell calcium homeostasis can also be remodeled during drug-induced cell differentiation. This indicates that defects of SERCA3 expression in cancer may be a widespread phenomenon, and show for the first time that ER calcium homeostatic defects are involved in shaping the lung cancer phenotype.

Results

SERCA expression in lung tumor cell lines and primary tumor tissue

Western immunoblot analysis performed on equal amounts of proteins obtained from total cell lysates from various non-small-cell lung carcinoma cell lines indicates that the SERCA expression pattern of these cells is heterogeneous. As shown in Fig. 1A, whereas all cell lines express the ubiquitous SERCA2 isoenzyme,

SERCA3 expression displays a marked heterogeneity: in cell lines such as HCC-827 or NCI-H441, SERCA3 protein expression is barely detectable or absent, but in other cell lines such as Calu-3, A-427, NCI-H23, A549, Cha-Go K-1, NCI-H1650, NCI-H358 or NCI-H460, SERCA3 protein is highly expressed and expression levels vary depending on cell line.

When investigated *in situ* by immunohistochemistry using a SERCA3-specific monoclonal antibody, SERCA3 expression could be detected in normal ciliated bronchial epithelium and, to a variable degree, also in primary human lung adenocarcinoma tissue. As shown in Fig. 1B, whereas bronchial epithelial cells were markedly labeled (++, photograph 2, black arrow) in a highly reproducible manner for SERCA3, staining in lung adenocarcinomas was heterogeneous and varied between weak (+/–, photograph 6), moderate (+, photographs 4 and 5) and marked (++, photograph 3). SERCA3 labeling in normal lung parenchyma (photograph 1) was readily detectable in alveolar macrophages (black arrowhead), and normal infiltrating lymphocytes displayed strong SERCA3 labeling (+++, white arrowheads) in normal (photograph 2) as well as tumor tissue (photographs 3–6).

These observations taken together show that SERCA3 expression is part of the normal differentiation program of lung epithelium, and that SERCA3 expres-

sion may be decreased or lost in lung adenocarcinoma. In addition, these observations indicate that heterogeneous SERCA3 expression in lung cancer cell lines *in vitro* reflects a pathophysiologically relevant phenomenon, because this is also observed in tumor tissue *in situ*.

Induction of SERCA expression by short chain fatty acids

When various non-small-cell lung carcinoma cell lines were subjected to treatments with short chain fatty acids and analogs, a selective induction of SERCA3 expression could be observed. As shown in Fig. 2, treatment with 5 mM phenylbutyrate for 5 days resulted in a marked and selective upregulation of the expression of the SERCA3 isoenzyme, whereas the expression of SERCA2 was not modified significantly or was decreased. SERCA3 induction could also be obtained by *n*-butyrate or *n*-valerate as well (not shown). As studied in the A549 lung adenocarcinoma cell line, induction of SERCA3 expression was concentration dependent in the low millimolar range and reached a plateau above 2 mM *n*-butyrate or 4-phenylbutyrate concentrations (Fig. 3A,B). SERCA3 expression reached a plateau of four- to six-fold without significant toxicity compared with untreated cells over a period of 4–7 days in cells treated with 3 mM butyrate or phenylbutyrate,




Fig. 1. SERCA expression in lung adenocarcinoma cell lines and lung tumor tissue. (A) Expression of SERCA2 and SERCA3 proteins in untreated lung carcinoma cell lines, detected with the IID8 (SERCA2) and the PLIM430 (SERCA3) monoclonal antibodies by western blotting. (B) Immunohistochemical detection of SERCA3 protein in normal lung parenchyma (photograph 1), bronchial epithelium (photograph 2) and grade 1 (photographs 3 and 4), grade 2 (photograph 6) and grade 3 (photograph 5) lung adenocarcinomas. Black arrows, normal bronchial epithelial cells; white arrowheads, normal lymphocytes; black arrowhead, alveolar macrophage (original magnification 40 \times).





Fig. 2. Selective induction of SERCA3 protein expression in various lung adenocarcinoma cell lines. The cells were treated with 5 mM phenylbutyrate (PB) for 5 days, and identical amounts of total cell protein lysates were analyzed using the IID8 (SERCA2-specific) and PLIM430 (SERCA3-specific) antibodies by western blotting. SERCA protein expression in phenylbutyrate treated cells (+) is compared with untreated controls (-).

Fig. 3. Dose–response relationship of SERCA3 induction. A549 cells were treated with various concentrations of butyrate (A) and phenylbutyrate (B), and SERCA expression was measured by semi-quantitative western blotting. An approximately six-fold induction of SERCA3 expression (black columns) is obtained in the low millimolar concentration range for both molecules, whereas the expression of SERCA2 (empty columns) decreases slightly (butyrate) or is not modified significantly (phenylbutyrate; $n = 3$).



respectively (Fig. S1A,B). Induction of SERCA3 expression could be obtained by other short chain fatty acid analogs as well. As shown in Fig. S2, when A549 cells were treated with short chain fatty acids of increasing chain length (from acetate to caproate), maximal induction could be obtained by butyrate and valerate. Among the tested branched chain fatty acid analogs, valproate was also active. Aryl-substituted analogs such as 3-phenylpropionate, 4-phenylbutyrate or 5-phenylvalerate also induced SERCA3 expression, with maximal activity observed in the case of phenylbutyrate.

In parallel with SERCA2 and SERCA3, the expression of gelsolin and p21^{CIP1/WAF1}, two general markers of lung cancer cell differentiation [48,49], has been investigated by western blotting in several lung adenocarcinoma cell lines treated with various short chain fatty acids for 5 days. As shown in Fig. 4, the selective induction of SERCA3 expression of A549, NCI-H358, NCI-H1650, Cha-Go K1 and HCC-827 cells with butyrate, valerate or phenylbutyrate was accompanied, in all investigated configurations, by increased gelsolin and p21^{CIP1/WAF1} expression. This indicates that the treatments induce phenotypic differentiation of the cells, and that the induction of SERCA3 expression is part of this process in all investigated model systems.

Induction of SERCA3 expression during the spontaneous differentiation of Calu-3 cells in post-confluent culture

Calu-3 lung adenocarcinoma cells have been shown to spontaneously undergo phenotypic differentiation towards a ciliated bronchial epithelial phenotype in post-confluent cultures [50]. The cells form a tight, polarized monolayer that displays trans-epithelial electrical resistance, barrier function and vectorial solute transport, as reflected by the formation of domes when grown on plastic, and express apical microvilli and tight junctional complexes [51]. Post-confluent Calu-3 cells are widely used in pharmacological research as an *in vitro* model for bronchial epithelium [52]. As shown in Fig. 5, the expression of SERCA3 is markedly increased in Calu-3 cells after confluency, whereas SERCA2 expression is not modified significantly. Interestingly, short chain fatty acid treatment of confluent Calu-3 led to a further increase of SERCA3 expression. Phenylbutyrate has already been shown to enhance CFTR protein expression in post-confluent, differentiated Calu-3 cells [52]. These data taken together indicate that the induction of SERCA3 expression takes place during the spontaneous differentiation of




Fig. 4. Expression of lung cancer cell differentiation markers by short chain fatty acids. Several lung adenocarcinoma cell lines (A549, NCI-H358, NCI-H1650, Cha-Go K1 and HCC-827) were treated with various short chain fatty acids (phenylbutyrate, 3 mM; valerate or butyrate, 5 mM) for 5 days, and SERCA expression as well as the expression of gelsolin (82 kDa) and p21^{CIP1/WAF1} (21 kDa) was observed by western blotting. Similarly to SERCA3, the expression of gelsolin and p21^{CIP1/WAF1} is increased in short chain fatty acid treated cells in all investigated drug/cell line configurations.

post-confluent Calu-3 cells, a process that can be further enhanced by phenylbutyrate.

Endoplasmic reticulum calcium storage in short chain fatty acid differentiated cells

In order to investigate ER calcium storage capacity in lung carcinoma cells, we established 11 clonal cell lines (BWV543 cells) derived from A549 lung adenocarcinoma that stably express the GCaMP2 recombinant fluorescent calcium probe in the cytosol and performed confocal microscope measurements of fluorescence on cells treated with 5 mM phenylbutyrate for 5 days and on untreated control cells. Treatment of all clones led to increased SERCA3 expression similarly to the

parental A549 cell line. ER calcium storage capacity was tested by confocal microscopy on two randomly selected clones (BWV543 clones 4 and 2.4). As shown in Fig. 6, when clone 4 cells incubated in the absence of extracellular calcium were treated with thapsigargin, a significant calcium release from the ER to the cytosol could be observed in control cells, whereas thapsigargin-induced calcium release in cells in which SERCA3 expression was induced by phenylbutyrate was significantly smaller. After the completion of thapsigargin-induced calcium release measured in the absence of extracellular calcium, 2 mM Ca²⁺ was added to the extracellular medium in order to observe capacitative calcium influx. Despite the marked decrease of thapsigargin-induced calcium release in phenylbutyrate treated cells, capacitative calcium influx was of essentially identical amplitude in untreated and phenylbutyrate treated cells. Identical results were obtained on BWV543 clone 2.4 cells as well.

Discussion

Because calcium uptake into the ER is performed exclusively by SERCA enzymes, SERCA function constitutes a key nodal point in cellular calcium homeostasis and signaling. SERCA activity is required for calcium accumulation in a resting cell for intra-ER calcium-dependent chaperone functions, as well as for calcium accumulation in the ER for inositol-1,4,5-trisphosphate-induced calcium release during cell activation. Moreover, because SERCA enzymes take up calcium from the cytosol even during an inositol-1,4,5-trisphosphate-induced calcium release event, they can shape the amplitude and the duration of calcium peaks, or modulate the spatiotemporal characteristics of calcium oscillations in the cytosol. SERCA activity can therefore critically modulate the state of activation of a cell, and can influence the type and the amplitude of the response given by a cell to calcium-dependent stimuli. This has important consequences for tumor biology: direct pharmacological SERCA inhibitors such as thapsigargin or 2,5-di-*tert*-butyl-1,4-benzohydroquinone are known tumor promoters, and SERCA2^{+/-} knockout is known to lead to squamous tumorigenesis with a long incubation time. In addition, sequencing of the SERCA2 and SERCA3 genes in various tumors including lung cancer and corresponding normal tissue suggests that inactivating mutations occur in these genes in a proportion of tumors [40,41], and it has thus been suggested that SERCA3 haploinsufficiency may predispose to cancer development or may promote tumor progression [41,42]. In this context the induction of SERCA3 expression, as observed




Fig. 5. Induction of SERCA3 expression during the spontaneous differentiation of post-confluent Calu-3 cells – superinduction by short chain fatty acids. Calu-3 lung adenocarcinoma cells were grown in pre-confluent conditions and in post-confluency, in the absence or the presence of various short chain fatty acids (5 mM) for 5 days, and SERCA expression was determined by western blotting. (A), (B) Photomicrographs of pre- and post-confluent cultures. Formation of ‘domes’ in post-confluent cultures (arrowheads) is indicative of trans-epithelial solute transport, a feature of a tight, differentiated epithelial monolayer. (C), (D) Cells at day 1 of post-confluency were allowed to undergo differentiation for 5 days in the absence or the presence of 5 mM butyrate, valerate or phenylbutyrate. SERCA2 and SERCA3 expression was then detected by western blotting and quantified. The selective induction of SERCA3 expression by post-confluent growth is further enhanced by short chain fatty acids. White bars, pre-confluent untreated cells; grey bars, post-confluent untreated cells; black bars, post-confluent, phenylbutyrate treated cells ($n = 3$).

in our work, may therefore exert anti-oncogenic effects in lung epithelial cells.

The observations presented in this work show, for the first time, that the expression of SERCA-type calcium pumps in lung cancer cells is dynamic, as various cell-differentiation-inducing stimuli led to the selective induction of the expression of the SERCA3 isoenzyme in all investigated cell lines. In addition, strong SERCA3 expression was found in fully differentiated bronchial epithelium. These data, taken together, indicate that SERCA3 expression is part of the differentiation program of normal bronchial epithelium and that this phenomenon can be recapitulated to various degrees during the differentiation of all investigated cell lines.

SERCA3 induction in A549 cells, a widely studied lung adenocarcinoma line, was associated with a decreased calcium release signal upon SERCA inhibition. The calcium affinity of SERCA3 is inferior (approximately 1.2 μM) to that of the co-expressed SERCA2b isoform (approximately 0.2 μM). The decreased calcium release signal observed in phenylbutyrate treated A549 cells cannot be attributed exclusively to SERCA3 induction due to the intercon-

nectedness of calcium homeostatic mechanisms that operate simultaneously in a cell. Decreased calcium release upon SERCA inhibition, however, is compatible with the notion of the formation of a leakier or lower-affinity, SERCA3-associated intracellular calcium pool, similar to that observed previously in butyrate treated KATO-III gastric carcinoma cells [18].

The normal lung contains several distinct and specialized epithelial cell types. In addition, lung carcinoma as a group consists of several histological tumor types that can display different degrees of histological differentiation. The ontogenic relationship between normal and tumoral cell types in the lung is not established in sufficient detail [1], and the molecular signaling mechanisms involved in the induction and the maintenance of tumoral phenotypes of increasing malignancy are not sufficiently known in the lung. Consequently, lung cancer is currently not amenable to successful pharmacological intervention. The observation that lung cancer cell calcium ER homeostasis is remodeled during differentiation is important for cancer biology because this indicates that lung adenocarcinoma cells may display calcium homeostatic defects




Fig. 6. Calcium release from SERCA-dependent intracellular calcium pools in A549 cells. A549 cells carrying the GCaMP2 fluorescent calcium probe (BWV543-4 cells) were treated or not with 5 mM phenylbutyrate for 5 days, and cytosolic calcium fluorescence was recorded by confocal fluorescence microscopy after complete inhibition of SERCA activity with 10 μ M thapsigargin in the absence of extracellular calcium. After completion of the thapsigargin-induced calcium release signal, extracellular medium was replenished with 2 mM Ca^{2+} and capacitative calcium influx was observed. Phenylbutyrate treatment leads to a significant decrease of calcium release from intracellular pools upon complete SERCA inhibition, whereas capacitative calcium influx remains similar. A total number of 36 treated and 48 untreated individual cells were recorded in four parallel experiments. Right panel: similarly to parental A549 cells, SERCA3 expression was induced by phenylbutyrate in A549 cells carrying the GCaMP2-carrying calcium probe.

similar to those observed in colon, gastric and breast carcinoma, as well as acute promyelocytic leukemia. In those systems the cross-talk of ER calcium homeostasis with other mechanisms of control of cell differentiation permits pharmacologically induced cell differentiation to be induced and potentiated and certain forms of resistance to differentiation-induction therapy to be overcome *in vitro* [44,46]. In addition, SERCA3 expression has also been shown to be a useful new immunohistochemical marker for the study of the state of differentiation and degree of malignancy of colon and breast tumors [44,45]. Data presented in this paper indicate, for the first time, that ER calcium biology is connected to cell differentiation also in pulmonary epithelium, and shows that ER calcium homeostasis presents defects in lung cancer cells, which can be overcome by pharmacologically induced cell differentiation. Although the precise correlation of the loss of SERCA3 expression with lung adenocarcinoma histological type, grade, molecular subtypes or clinical parameters requires further work, the demonstration of calcium homeostatic anomalies in lung cancer may help identify new calcium-dependent targets for the therapy

of the disease. In addition, our observations suggest that SERCA3 may prove useful for the immunohistochemical analysis of lung tumors and for the study of bronchic epithelial differentiation.

Materials and methods

Chemicals

n-Butyric and *n*-valeric, 4-phenylbutyric acid, as well as sodium acetate, sodium caproate, sodium valproate, isobutyric, 4-methylvaleric as well as phenylacetic, 3-phenylpropionic and 5-phenylvaleric acids were from Sigma-Aldrich (Saint-Quentin Fallavier, France). Free acids were dissolved to 300 mM final concentration in 300 mM sterile sodium bicarbonate solution, filtered through 0.2 μ m membranes and stored at -25°C . Sodium salts were dissolved at 300 mM concentration in sterile water and filtered through 0.2 μ m membranes.

Cell lines

The Calu-3, NCI-H23, NCI-H358, NCI-H441, NCI-H460, NCI-H596, NCI-H1299 and NCI-H1650 as well as the HCC-827 cell lines were obtained from ATCC (ATCC-LGC Standards Sarl, Molsheim, France). A549 and A427 cells were from DSMZ (Braunschweig, Germany), and Cha-Go K1 was purchased from ECACC (Porton Down, UK). The various cell lines were cultured according to the instructions of the cell line depository of origin. RPMI-based media contained Ultraglutamine-I (Lonza, Verviers, Belgium) in addition to 2 mM glutamine.

Treatments

Exponentially growing cells were trypsinized and plated into 60 cm^2 cell culture grade Petri dishes in new medium. When cultures reached 40–60% confluency as examined by light microscopy, medium was renewed and drugs were added from concentrated stock solutions. Following treatments as indicated in the figures, cells were washed with ice-cold 150 mM NaCl and precipitated with 5% trichloroacetic acid. The trichloroacetic acid pellet was quantified and dissolved in modified Laemmli-type sample buffer exactly as described earlier [17], and 50 μ g total cellular protein was deposited for SDS/polyacrylamide gel electrophoresis per well [17]. Following electrophoresis and transfer to nitrocellulose membranes, equal loading of samples was controlled prior immunoblotting by Ponceau red staining of total protein deposited on the membranes. Ponceau red stained blots were scanned and equal protein loading was controlled by densitometry using the SCIONIMAGE software (Scion Corporation, Frederick, MD, USA) as described earlier in detail [17].

Western blotting

The SERCA3-specific PLIM430 mouse IgG1 monoclonal antibody was used for western immunoblotting as hybridoma supernatant as described earlier in detail [17,18,44]. Purified mouse anti-gelsolin antibody (Cat # 610412; BD Transduction Laboratories, San José, CA, USA) (dissolved in NaCl/Tris milk at $0.05 \mu\text{g}\cdot\text{mL}^{-1}$ final concentration) and the mouse monoclonal anti-p21 antibody SC-71811 (Santa Cruz Biotechnologies, CA, USA) (dissolved in NaCl/Tris milk at $0.2 \mu\text{g}\cdot\text{mL}^{-1}$ final concentration) were used on heat treated samples (100°C for 12 min) run in 8% polyacrylamide gels. Antibodies were revealed with an anti-mouse Ig horseradish peroxidase conjugate (Jackson Immuno-Research, Newmarket, Suffolk, UK) and the enhanced chemiluminescence system of Amersham (Courtaboeuf, France) as described earlier [17,18,44]. Non-saturated luminograms were scanned and densitometric analysis was done with the SCIONIMAGE software [11–15].

Immunohistochemistry

Formalin fixed paraffin embedded lung cancer tissue microarray slides were purchased from BioChain Institute, Hayward, CA, USA (T8235724, T8235732, Z7020066, Z7020067; CliniSciences, Montrouge, France), and from US Biomax, Rockville, MD, USA (BC04015, BC041114, BC041115, BC04119b, LC20810; Euromedex, Souffelweyersheim, France). Staining for SERCA3 was performed using the clone 2H3 monoclonal mouse anti-SERCA3 antibody (IgG2a kappa) from Abnova (Tebu-bio, Le Perray en Yvelines, France) raised against a GST-tagged peptide sequence encompassing the 501–621 amino acid fragment of SERCA3, at $3 \mu\text{g}\cdot\text{mL}^{-1}$ final concentration diluted in Dako REAL antibody diluent (Dako France SAS, Trappes, France) as described in [45]. Briefly, after inhibition of endogenous peroxidase and antigen retrieval of deparaffinized sections by a *tris*-hydroxymethyl-aminomethane-based reagent (CC1 solution; Ventana Medical Systems, Illkirch, France) at $95\text{--}100^\circ\text{C}$ for 12 min, slides were incubated for 30 min at 37°C with the antibody, and staining was revealed using the Ventana I-View Biotin-Ig-streptavidin-biotin-horseradish peroxidase system with blocking of endogenous biotin activity and copper enhancement, according to the instructions of the manufacturer. As negative control, isotype matched irrelevant antibody was used at the same concentration, and this gave no staining. As internal positive control, lymphocytes and vascular endothelial cells present in the samples and which express high levels of SERCA3 [17,24] were used. Slides were counterstained with hematoxylin and bluing agent (Ventana). SERCA3 immunostaining of the samples was evaluated semi-quantitatively using a four-grade scale (0 to 3+). Photomicrographs were taken with a Zeiss Axio Scope.A1 microscope equipped with a Zeiss N-Achroplan $40\times/0.65$ objective and an AxioCam

ICCl camera, using the AXIOVISION 4.8.2 software. Tissue microarray slides were also processed using the ImmPRESS biotin-free polymerized enzyme staining system (Vector Laboratories, Clinisciences SAS, Nanterre, France) according to the instructions of the manufacturer, and SERCA3 staining with this system gave identical results.

Confocal fluorescent microscope calcium measurements

In order to generate the SB-CAG-GCaMP2 fluorescent calcium probe, a DNA fragment encoding GCaMP2 was first generated by PCR using the pN1-GCaMP2 plasmid, a generous gift of Junichi Nakai, RIKEN Brain Science Institute, Saitama, Japan [53]. Primer sequences were as follows: forward, 5'-CTACCGGTCTCGCCACCAATG-3'; reverse, 5'-AGATCTCCGCTCACTTCGCTGTC-3'. The forward primer adds an *AgeI* restriction site, the reverse primer adds a *BglII* restriction site. The complete GCaMP2 DNA fragment was cloned via *AgeI*(5')/*BglII*(3') into an SB-CAG-AmaraGFP vector from which Amara GFP was excised with the same enzyme pair. The GCaMP2 fragment is framed in the resulting vector by two left inverted repeat–direct repeat regions, which are the recognition motifs of *Sleeping Beauty* (SB) transposase [54–57], and possesses a cytomegalovirus early enhancer/chicken β actin (CAG) [58] promoter. The efficiency of such a symmetrical transposon was proved by T. Orbán (personal communication). This construct was used for stable transfection using the SB transposon delivery system as described earlier in detail [54–57,59]. For the SB transposase, we used an enhanced version of the enzyme having an approximately 100-fold higher activity than the originally reconstructed transposase [59,60]. For transfection of exponentially growing A549 cells, the FuGENE[®] 6 (Roche Applied Science, Rotkreuz, Switzerland, <http://www.roche-applied-science.com>) reagent was used according to the instructions of the manufacturer. Cells were cotransfected with transposon and transposase plasmids in a 10:1 ratio to avoid overproduction inhibition of the transposase [56,57]. Following transfections, cells were cultured for 3 days and then resuspended by trypsinization and sorted for GFP fluorescence using the Aria High Speed Cell Sorter of Becton-Dickinson (San José, CA, USA, <http://www.bdbiosciences.com>). The sorted cell population (containing approximately 60% GFP positive cells) was further cultured in bulk for 4 days, trypsinized and single cell cloned by limiting dilution in 96-well plates. Individual GCaMP2 expressing clones were identified by GFP fluorescence using an inverted fluorescence microscope and expanded.

SB-CAG-GCaMP2-carrying A549 clones were grown to 20–30% confluency in eight-chamber Lab-Tek glass bottom chamber slides and treated with 5 mM phenylbutyrate for 5 days. At the end of treatment, the cells were washed with

calcium-free Hanks' solution. Cells were then observed with an Olympus IX-81/FV500 laser scanning confocal microscope using an Olympus PLAPO 60× (1.4) oil immersion objective. For GCaMP2 imaging cells were excited with the 488 nm laser line and emission was collected between 505 and 535 nm. Under these conditions emitted fluorescence is proportional to the cytosolic free calcium concentration [53].

Calcium signal measurements were carried out in Hanks' balanced salt solution supplemented with 20 mM Hepes (pH 7.4) and 0.9 mM MgCl₂. This medium was supplemented with 100 μM CaCl₂ and 100 μM EGTA for the measurement of thapsigargin-induced calcium release from the ER. Under these conditions free calcium concentration in the medium is approximately 4 μM, and no calcium influx is observed. Capacitative calcium influx was thereafter induced by the addition of 2 mM free CaCl₂ to the medium. Time lapse sequences of cellular fluorescence were recorded and images were analyzed with the FLUOVIEW TIEMPO (v4.3; Olympus, <http://www.olympusmicro.com>) time course software. In order to compensate for variability of fluorophore expression levels in individual cells, fluorescence values are expressed as F/F_0 . A total of 36 untreated and 48 treated cells were recorded individually in four parallel experiments.

Statistical analysis

Data are presented as the mean ± SEM and correspond to at least three independent experiments. Statistical analysis was done using Student's paired *t* test.

Acknowledgements

We express special thanks to Professor Neville Crawford (Department of Biochemistry, Hunterian Institute, London, UK) for giving us the PLIM430 hybridoma, and to Dr Tamás Orbán (Membrane Research Group, Hungarian Academy of Sciences, Hungary) for his contribution regarding the use of the transposon system used in this work. The excellent technical help of Mr Patrice Castagnet (Service d'Anatomie et Cytologie Pathologiques, Hôpital Lariboisière, Paris, France) is gratefully acknowledged. This work was supported by INSERM, the Association pour la Recherche sur le Cancer, France, and by the Hungarian Academy of Sciences grants CK80283 and K101064 (EA). This work is dedicated to the memory of Casper and Rose.

References

- 1 Travis WD (2004) World Health Organization, International Agency for Research on Cancer, International Association for the Study of Lung Cancer, International Academy of Pathology: Pathology and Genetics of Tumours of the Lung, Pleura, Thymus, and Heart. IARC Press, Lyon.
- 2 Youlten DR, Cramb SM & Baade PD (2008) The international epidemiology of lung cancer: geographical distribution and secular trends. *J Thorac Oncol* **3**, 819–831.
- 3 Berridge MJ (2009) Inositol trisphosphate and calcium signalling mechanisms. *Biochim Biophys Acta* **1793**, 933–940.
- 4 Berridge MJ (2002) The endoplasmic reticulum: a multifunctional signaling organelle. *Cell Calcium* **32**, 235–249.
- 5 Coe H & Michalak M (2009) Calcium binding chaperones of the endoplasmic reticulum. *Gen Physiol Biophys* **28**, F96–F103.
- 6 Malhotra JD & Kaufman RJ (2007) The endoplasmic reticulum and the unfolded protein response. *Semin Cell Dev Biol* **18**, 716–731.
- 7 Burdakov D, Petersen OH & Verkhratsky A (2005) Intraluminal calcium as a primary regulator of endoplasmic reticulum function. *Cell Calcium* **38**, 303–310.
- 8 Bánhegyi G, Baumeister P, Benedetti A, Dong D, Fu Y, Lee AS, Li J, Mao C, Margittai E, Ni M *et al.* (2007) Endoplasmic reticulum stress. *Ann N Y Acad Sci* **1113**, 58–71.
- 9 Lai E, Teodoro T & Volchuk A (2007) Endoplasmic reticulum stress: signaling the unfolded protein response. *Physiology* **22**, 193–201.
- 10 Wuytack F, Raeymaekers L & Missiaen L (2002) Molecular physiology of the SERCA and SPCA pumps. *Cell Calcium* **32**, 279–305.
- 11 Vangheluwe P, Sepulveda MR, Missiaen L, Raeymaekers L, Wuytack F & Vanoevelen J (2009) Intracellular Ca²⁺- and Mn²⁺-transport ATPases. *Chem Rev* **109**, 4733–4759.
- 12 Vandecaetsbeek I, Vangheluwe P, Raeymaekers L, Wuytack F & Vanoevelen J (2011) The Ca²⁺ pumps of the endoplasmic reticulum and Golgi apparatus. *Cold Spring Harb Perspect Biol* **3**, a004184. doi:10.1101/cshperspect.a004184.
- 13 Bobe R, Bredoux R, Corvazier E, Lacabaratz-Porret C, Martin V, Kovács T & Enouf J (2005) How many Ca²⁺-ATPase isoforms are expressed in a cell type? A growing family of membrane proteins illustrated by studies in platelets. *Platelets* **16**, 133–150.
- 14 Dally S, Corvazier E, Bredoux R, Bobe R & Enouf J (2010) Multiple and diverse coexpression, location, and regulation of additional SERCA2 and SERCA3 isoforms in nonfailing and failing human heart. *J Mol Cell Cardiol* **48**, 633–644.
- 15 Hovnanian A (2007) SERCA pumps and human diseases. *Subcell Biochem* **45**, 337–363.
- 16 Arbabian A, Brouland JP, Gélébart P, Kovács T, Bobe R, Enouf J & Papp B (2011) Endoplasmic

- reticulum calcium pumps and cancer. *BioFactors* **37**, 139–149.
- 17 Dellis O, Arbabian A, Brouland JP, Kovács T, Rowe M, Chomienne C, Joab I & Papp B (2009) Modulation of B-cell endoplasmic reticulum calcium homeostasis by Epstein–Barr virus latent membrane protein-1. *Mol Cancer* **8**, 59.
 - 18 Gélébart P, Kovács T, Brouland JP, van Gorp R, Grossmann J, Rivard N, Panis Y, Martin V, Bredoux R, Enouf J *et al.* (2002) Expression of endomembrane calcium pumps in colon and gastric cancer cells. Induction of SERCA3 expression during differentiation. *J Biol Chem* **277**, 26310–26320.
 - 19 Launay S, Bobe R, Lacabartz-Porret C, Bredoux R, Kovács T, Enouf J & Papp B (1997) Modulation of endoplasmic reticulum calcium pump expression during T lymphocyte activation. *J Biol Chem* **272**, 10746–10750.
 - 20 Launay S, Gianni M, Kovács T, Bredoux R, Bruel A, Gélébart P, Zasadowski F, Chomienne C, Enouf J & Papp B (1999) Lineage-specific modulation of calcium pump expression during myeloid differentiation. *Blood* **93**, 4395–4405.
 - 21 Papp B, Enyedi A, Kovács T, Sarkadi B, Wuytack F, Thastrup O, Gárdos G, Bredoux R, Lèvy-Toledano S & Enouf J (1991) Demonstration of two forms of calcium pumps by thapsigargin inhibition and radioimmunoblotting in platelet membrane vesicles. *J Biol Chem* **266**, 14593–14596.
 - 22 Papp B, Enyedi A, Pászty K, Kovács T, Sarkadi B, Gárdos G, Magnier C, Wuytack F & Enouf J (1992) Simultaneous presence of two distinct endoplasmic-reticulum-type calcium-pump isoforms in human cells. Characterization by radio-immunoblotting and inhibition by 2,5-di-(*t*-butyl)-1,4-benzohydroquinone. *Biochem J* **288**, 297–302.
 - 23 Wuytack F, Papp B, Verboomen H, Raeymaekers L, Dode L, Bobe R, Enouf J, Bokkala S, Authi KS & Casteels R (1994) A sarco/endoplasmic reticulum Ca^{2+} -ATPase 3-type Ca^{2+} pump is expressed in platelets, in lymphoid cells, and in mast cells. *J Biol Chem* **269**, 1410–1416.
 - 24 Hadri L, Ozog A, Soncin F & Lompré AM (2002) Basal transcription of the mouse sarco(endo)plasmic reticulum Ca^{2+} -ATPase type 3 gene in endothelial cells is controlled by Ets-1 and Sp1. *J Biol Chem* **277**, 36471–36478.
 - 25 Diederichs F (2008) Ion homeostasis and the functional roles of SERCA reactions in stimulus-secretion coupling of the pancreatic beta-cell: a mathematical simulation. *Biophys Chem* **134**, 119–143.
 - 26 Váradi A, Molnár E, Ostenson CG & Ashcroft SJ (1996) Isoforms of endoplasmic reticulum Ca^{2+} -ATPase are differentially expressed in normal and diabetic islets of Langerhans. *Biochem J* **319**, 521–527.
 - 27 Bobe R, Bredoux R, Corvazier E, Andersen JP, Clausen JD, Dode L, Kovács T & Enouf J (2004) Identification, expression, function and localization of a novel (sixth) isoform of the human sarco/endoplasmic reticulum Ca^{2+} ATPase 3 gene. *J Biol Chem* **279**, 24297–24306.
 - 28 Dode L, Vilsen B, Van Baelen K, Wuytack F, Clausen JD & Andersen JP (2002) Dissection of the functional differences between sarco(endo)plasmic reticulum Ca^{2+} -ATPase (SERCA) 1 and 3 isoforms by steady-state and transient kinetic analyses. *J Biol Chem* **277**, 45579–45591.
 - 29 Poch E, Leach S, Snape S, Cacic T, MacLennan DH & Lytton J (1998) Functional characterization of alternatively spliced human SERCA3 transcripts. *Am J Physiol* **275**, C1449–1458.
 - 30 Wuytack F, Dode L, Baba-Aissa F & Raeymaekers L (1995) The SERCA3-type of organellar Ca^{2+} pumps. *Biosci Rep* **15**, 299–306.
 - 31 Lytton J, Westlin M, Burk SE, Shull GE & MacLennan DH (1992) Functional comparisons between isoforms of the sarcoplasmic or endoplasmic reticulum family of calcium pumps. *J Biol Chem* **267**, 14483–14489.
 - 32 Chandrasekera PC, Kargacin ME, Deans JP & Lytton J (2009) Determination of apparent calcium affinity for endogenously expressed human sarco(endo)plasmic reticulum calcium-ATPase isoform SERCA3. *Am J Physiol Cell Physiol* **296**, C1105–1114.
 - 33 Lipskaia L, Hulot JS & Lompré AM (2009) Role of sarco/endoplasmic reticulum calcium content and calcium ATPase activity in the control of cell growth and proliferation. *Pflugers Arch* **457**, 673–685.
 - 34 Apáti A, Jánossy J, Brózik A, Bauer PI & Magócsi M (2003) Calcium induces cell survival and proliferation through the activation of the MAPK pathway in a human hormone-dependent leukemia cell line, TF-1. *J Biol Chem* **278**, 9235–9243.
 - 35 Brini M & Carafoli E (2009) Calcium pumps in health and disease. *Physiol Rev* **89**, 1341–1378.
 - 36 Gill DL, Waldron RT, Rys-Sikora KE, Ufret-Vincenty CA, Graber MN, Favre CJ & Alfonso A (1996) Calcium pools, calcium entry, and cell growth. *Biosci Rep* **16**, 139–157.
 - 37 Sakai A & Teshima R (2001) 2,5-Di-*tert*-butyl-1,4-hydroquinone enhances cell transformation accompanied by an increase in intracellular free calcium ion concentration. *Cancer Lett* **168**, 183–190.
 - 38 Hakii H, Fujiki H, Suganuma M, Nakayasu M, Tahira T, Sugimura T, Scheuer PJ & Christensen SB (1986) Thapsigargin, a histamine secretagogue, is a non-12-O-tetradecanoylphorbol-13-acetate (TPA) type tumor promoter in two-stage mouse skin carcinogenesis. *J Cancer Res Clin Oncol* **111**, 177–181.

- 39 Prasad V, Okunade GW, Miller ML & Shull GE (2004) Phenotypes of SERCA and PMCA knockout mice. *Biochem Biophys Res Commun* **322**, 1192–1203.
- 40 Korošec B, Glavac D, Rott T & Ravnik-Glavac M (2006) Alterations in the ATP2A2 gene in correlation with colon and lung cancer. *Cancer Genet Cytogenet* **171**, 105–111.
- 41 Korošec B, Glavac D, Volavsek M & Ravnik-Glavac M (2008) Alterations in genes encoding sarcoplasmic-endoplasmic reticulum Ca²⁺ pumps in association with head and neck squamous cell carcinoma. *Cancer Genet Cytogenet* **181**, 112–118.
- 42 Korošec B, Glavac D, Volavsek M & Ravnik-Glavac M (2009) ATP2A3 gene is involved in cancer susceptibility. *Cancer Genet Cytogenet* **188**, 88–94.
- 43 Liu LH, Boivin GP, Prasad V, Periasamy M & Shull GE (2001) Squamous cell tumors in mice heterozygous for a null allele of Atp2a2, encoding the sarco(endoplasmic reticulum Ca²⁺-ATPase isoform 2 Ca²⁺ pump. *J Biol Chem* **276**, 26737–26740.
- 44 Brouland JP, Gélébart P, Kovács T, Enouf J, Grossmann J & Papp B (2005) The loss of sarco/endoplasmic reticulum calcium transport ATPase 3 expression is an early event during the multistep process of colon carcinogenesis. *Am J Pathol* **167**, 233–242.
- 45 Papp B & Brouland JP (2011) Altered endoplasmic reticulum calcium pump expression during breast tumorigenesis. *Breast Cancer (Auckl)* **5**, 163–174.
- 46 Launay S, Gianni M, Diomedede L, Machesky LM, Enouf J & Papp B (2003) Enhancement of ATRA-induced cell differentiation by inhibition of calcium accumulation into the endoplasmic reticulum: cross-talk between RAR alpha and calcium-dependent signaling. *Blood* **101**, 3220–3228.
- 47 Papp B, Brouland JP, Gélébart P, Kovács T & Chomienne C (2004) Endoplasmic reticulum calcium transport ATPase expression during differentiation of colon cancer and leukaemia cells. *Biochem Biophys Res Commun* **322**, 1223–1236.
- 48 Chang TH & Szabó E (2000) Induction of differentiation and apoptosis by ligands of peroxisome proliferator-activated receptor gamma in non-small cell lung cancer. *Cancer Res* **60**, 1129–1138.
- 49 Chang TH & Szabó E (2002) Enhanced growth inhibition by combination differentiation therapy with ligands of peroxisome proliferator-activated receptor-gamma and inhibitors of histone deacetylase in adenocarcinoma of the lung. *Clin Cancer Res* **8**, 1206–1212.
- 50 Foster KA, Avery ML, Yazdaniyan M & Audus KL (2000) Characterization of the Calu-3 cell line as a tool to screen pulmonary drug delivery. *Int J Pharm* **208**, 1–11.
- 51 Mathia NR, Timoszyk J, Stetsko PI, Megill JR, Smith RL & Wall DA (2002) Permeability characteristics of calu-3 human bronchial epithelial cells: *in vitro*–*in vivo* correlation to predict lung absorption in rats. *J Drug Target* **10**, 31–40.
- 52 Löffing J, Moyer BD, Reynolds D & Stanton BA (1999) PBA increases CFTR expression but at high doses inhibits Cl[−] secretion in Calu-3 airway epithelial cells. *Am J Physiol* **277**, L700–708.
- 53 Nakai J, Ohkura M & Imoto K (2001) A high signal-to-noise Ca²⁺ probe composed of a single green fluorescent protein. *Nat Biotechnol* **19**, 137–141.
- 54 Ivics Z, Hackett PB, Plasterk RH & Izsvák Z (1997) Molecular reconstruction of Sleeping Beauty, a Tc1-like transposon from fish, and its transposition in human cells. *Cell* **91**, 501–510.
- 55 Ivics Z & Izsvák Z (2006) Transposons for gene therapy! *Curr Gene Ther* **6**, 593–607.
- 56 Izsvák Z & Ivics Z (2004) Sleeping beauty transposition: biology and applications for molecular therapy. *Mol Ther* **9**, 147–156.
- 57 Izsvák Z, Ivics Z & Plasterk RH (2002) Sleeping Beauty, a wide host-range transposon vector for genetic transformation in vertebrates. *J Mol Biol* **302**, 93–102.
- 58 Alexopoulou AN, Couchman JR & Whiteford JR (2008) The CMV early enhancer/chicken beta actin (CAG) promoter can be used to drive transgene expression during the differentiation of murine embryonic stem cells into vascular progenitors. *BMC Cell Biol* **9**, 2.
- 59 Mates L, Chuah MK, Belay E, Jerchow B, Manoj N, Acosta-Sanchez A, Grzela DP, Schmitt A, Becker K, Mátrai J *et al.* (2009) Molecular evolution of a novel hyperactive Sleeping Beauty transposase enables robust stable gene transfer in vertebrates. *Nat Genet* **41**, 753–761.
- 60 Orbán TI, Apáti A, Németh A, Varga N, Krizsik V, Schamberger A, Szebenyi K, Erdei Z, Várady G, Karázi E *et al.* (2009) Applying a ‘double-feature’ promoter to identify cardiomyocytes differentiated from human embryonic stem cells following transposon-based gene delivery. *Stem Cells* **27**, 1077–1087.

Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web site:

Fig. S1. Time course of SERCA2 and SERCA3 expression in short chain fatty acid treated A549 cells.

Fig. S2. Induction of SERCA3 expression in A549 cells by various short chain fatty analogs.